

Chapter 6

Home Dialysis



2016
ANZDATA Registry
39th Annual Report

Data to 31-Dec-2015

Introduction

At the end of 2015 30% of Australian and 48% of New Zealand dialysis patients were dialysing at home. This chapter reports the incidence, prevalence and outcomes of home dialysis, defined simply as the combination of peritoneal dialysis (PD) and home haemodialysis (HHD).

New Patients

In this section “incident home dialysis patients” are defined as those who commenced home dialysis for the first time (ie having never received home dialysis before), including those who had previously received facility haemodialysis or a kidney transplant.

Table 6.1 presents the number of incident patients (per million population) over time. There has been a slow increase in the number of patients commencing home dialysis over the last 20 years, predominantly driven by population growth.

Figure 6.1 shows the age distribution of incident home dialysis patients in 2015. The majority of incident home dialysis patients commence peritoneal dialysis, and the majority are aged 45-74. Figure 6.2 presents the same data per million population.

Table 6.1 Number (pmp) of Incident Home Dialysis Patients 1996-2015

Year	Australia	New Zealand
1996	829 (45)	252 (68)
1997	810 (44)	274 (72)
1998	870 (47)	302 (79)
1999	906 (48)	306 (80)
2000	934 (49)	305 (79)
2001	975 (51)	326 (84)
2002	915 (47)	347 (88)
2003	943 (48)	317 (79)
2004	888 (45)	332 (81)
2005	974 (48)	318 (77)
2006	1183 (58)	351 (84)
2007	1076 (52)	289 (68)
2008	1163 (55)	333 (78)
2009	1070 (49)	365 (85)
2010	922 (42)	366 (84)
2011	998 (45)	312 (71)
2012	1226 (54)	345 (78)
2013	1179 (51)	381 (86)
2014	1303 (56)	344 (76)
2015	1181 (50)	350 (76)

Figure 6.1.1




Figure 6.1.2






Figure 6.2.1**Figure 6.2.2**

The primary renal disease of incident home dialysis patients is shown in table 6.2. The distribution of these diseases is similar to the broader incident renal replacement therapy population (see chapter 1).

Table 6.2. Primary Disease (%) of Incident Home Dialysis Patients 2015

Primary Renal Disease	Australia	New Zealand
GN	285 (24)	81 (23)
Analgesic	10 (1)	0 (0)
Polycystic	81 (7)	12 (3)
Reflux	23 (2)	9 (3)
Hypertension	154 (13)	30 (9)
Diabetes	384 (33)	165 (47)
Other	146 (12)	39 (11)
Uncertain	57 (5)	10 (3)
Not reported	41 (3)	4 (1)
Total	1181 (100)	350 (100)

Figure 6.3 presents the cumulative incidence of patients commencing home dialysis over 2006-2015. The data are censored at transplantation, and death is handled as a competing risk. PD uptake is rapid, both as the first dialysis modality and within 6 months of starting dialysis, whereas HHD uptake is more gradual. Both modalities are taken up more rapidly in New Zealand than in Australia. Uptake also varies by state (figure 6.4), age (figure 6.5) and gender (figure 6.6).

Figure 6.3




Figure 6.4




Figure 6.5.1




Figure 6.5.2





Figure 6.6



Prevalent Patients

Figure 6.7 shows the number of patients dialysing in Australia and New Zealand at the end of 1996-2015. Although numbers of home dialysis patients are growing, this growth is substantially outpaced by the growth in the numbers of facility haemodialysis patients.

Figure 6.8 presents the age distribution of prevalent home dialysis patients, and figure 6.9 shows prevalence per million population. Figure 6.10 shows the geographical distribution of home dialysis patients in Australia at the end of 2015; mapping data are courtesy of the Australian Bureau of Statistics.

Figure 6.7




Figure 6.8.1




Figure 6.8.2




Figure 6.9.1




Figure 6.9.2





Figure 6.10

Prevalent home dialysis patients 2015
By statistical subdivision



Figures 6.11-6.12 show trends in the numbers of home dialysis patients by year and state (figure 6.11) and age (figure 6.12). The total number of home dialysis patients in each state at the end of 2015 is shown in table 6.3. These numbers should be interpreted in the broader context of the overall growth in dialysis prevalence (see figure 6.7 and chapter 2).

Figure 6.11




Figure 6.12.1




Figure 6.12.2




Table 6.3. Home Dialysis Patient Numbers by Country and State 2015

State	PD	Home HD	Other HD	Total
QLD	502	284	1583	2369
NSW/ACT	996	489	2700	4185
VIC	600	214	2207	3021
TAS	48	26	167	241
SA	118	34	609	761
NT	28	37	549	614
WA	222	81	967	1270
NZ	791	482	1401	2674
Total	3305	1647	10183	15135

The use of home dialysis varies substantially between treating hospitals. Of patients receiving maintenance dialysis at the end of 2015, the proportion who were dialysing at home varied from 0-100% in Australia and 31-88% in New Zealand (figure 6.13).

Figure 6.13.1





Figure 6.13.2



Outcomes of incident home dialysis patients

The outcomes of incident home dialysis patients are shown in table 6.4. In patients who commenced home dialysis for the first time during 2004-2015, the most common reasons for completion of home dialysis were transfer to facility haemodialysis and death.



Table 6.4. Outcome of Incident Home Dialysis Patients 2004-2015

Outcome	Australia	New Zealand
Transferred to facility haemodialysis	4702 (36%)	1316 (32%)
Died	2682 (20%)	1189 (29%)
Transplanted	2498 (19%)	539 (13%)
Lost to follow-up	26 (<1%)	6 (<1%)
Recovered native kidney function	236 (2%)	58 (1%)
Remained on home dialysis 31 Dec 2015	3019 (23%)	978 (24%)
Total	13163 (100%)	4086 (100%)



Patient and Technique Survival

The following pages present three outcomes of patients who commenced home dialysis for the first time over 2004-2015: (1) patient survival, censored at transplantation but not at dialysis modality change; (2) technique survival, censored at transplantation, and with technique failure defined as a change to facility haemodialysis for ≥ 30 days or death; and (3) death-censored technique survival, censored at transplantation and death, and with technique failure defined as a change to facility haemodialysis for ≥ 30 days. Home dialysis technique survival can be interpreted as how long patients who have commenced home dialysis are able to keep dialysing at home.

Each outcome is presented stratified by era (figures 6.14-6.16 and tables 6.5-6.7) and by age group (figures 6.17-6.19 and tables 6.8-6.10). Recent eras have seen an improvement in patient survival, and a minor improvement in technique (but not death-censored technique) survival. Age is strongly associated with patient and technique survival, but not death-censored technique survival, indicating that the association with technique survival is due to differences in patient survival rather than differences in the rate of transfer to facility haemodialysis.

Figure 6.14.1**Figure 6.14.2****Table 6.5. Incident Home Dialysis Patient Survival 2004-2015**

Country	Era	6 months	1 year	3 years	5 years
Australia	2004-2006 (n=3045)	95 (94, 95)	89 (88, 90)	66 (65, 68)	51 (49, 53)
	2007-2009 (n=3309)	96 (95, 96)	91 (90, 92)	71 (69, 73)	55 (53, 57)
	2010-2012 (n=3146)	97 (96, 97)	93 (92, 94)	75 (73, 76)	57 (55, 60)
	2013-2015 (n=3663)	98 (97, 98)	94 (93, 95)	-	-
New Zealand	2004-2006 (n=1001)	95 (93, 96)	89 (87, 91)	66 (63, 69)	45 (41, 48)
	2007-2009 (n=987)	96 (95, 97)	90 (88, 92)	69 (66, 72)	49 (46, 52)
	2010-2012 (n=1023)	97 (96, 98)	93 (91, 94)	69 (66, 72)	46 (42, 50)
	2013-2015 (n=1075)	97 (96, 98)	93 (91, 94)	-	-

Figure 6.15.1**Figure 6.15.2****Table 6.6. Incident Home Dialysis Technique Survival by Era 2004-2015**

Country	Era	6 months	1 year	3 years	5 years
Australia	2004-2006 (n=3045)	85 (83, 86)	73 (71, 74)	38 (36, 40)	20 (19, 22)
	2007-2009 (n=3309)	84 (83, 85)	72 (71, 74)	39 (37, 41)	22 (20, 23)
	2010-2012 (n=3146)	86 (84, 87)	75 (73, 76)	44 (42, 46)	23 (21, 26)
	2013-2015 (n=3663)	89 (88, 90)	80 (78, 81)	-	-
New Zealand	2004-2006 (n=1001)	88 (86, 90)	76 (74, 79)	43 (40, 46)	22 (19, 25)
	2007-2009 (n=987)	89 (87, 91)	79 (77, 82)	46 (42, 49)	25 (22, 28)
	2010-2012 (n=1023)	89 (87, 91)	80 (78, 83)	45 (42, 49)	21 (18, 24)
	2013-2015 (n=1075)	87 (85, 89)	78 (75, 81)	-	-

Figure 6.16.1




Figure 6.16.2








Table 6.7. Incident Home Dialysis Death-Censored Technique Survival by Era 2004-2015

Country	Era	6 months	1 year	3 years	5 years
Australia	2004-2006 (n=3045)	89 (88, 90)	81 (79, 82)	56 (53, 58)	41 (38, 43)
	2007-2009 (n=3309)	88 (87, 89)	79 (78, 81)	54 (52, 56)	41 (39, 43)
	2010-2012 (n=3146)	88 (87, 89)	80 (79, 81)	58 (56, 60)	42 (39, 45)
	2013-2015 (n=3663)	91 (90, 92)	85 (83, 86)	-	-
New Zealand	2004-2006 (n=1001)	92 (90, 94)	86 (83, 88)	65 (61, 68)	51 (46, 55)
	2007-2009 (n=987)	93 (91, 94)	87 (84, 89)	65 (62, 69)	50 (46, 54)
	2010-2012 (n=1023)	91 (89, 93)	86 (84, 88)	66 (63, 70)	47 (42, 52)
	2013-2015 (n=1075)	90 (88, 92)	84 (81, 86)	-	-

Figure 6.17.1**Figure 6.17.2****Table 6.8. Incident Home Dialysis Patient Survival by Age 2004-2015**

Country	Age	6 months	1 year	3 years	5 years
Australia	<40 (n=2035)	98 (98, 99)	97 (96, 97)	90 (88, 91)	85 (83, 87)
	40-59 (n=4675)	97 (97, 98)	95 (94, 95)	80 (79, 82)	67 (65, 69)
	60-74 (n=4567)	96 (95, 96)	90 (89, 91)	66 (64, 67)	47 (45, 49)
	≥75 (n=1886)	92 (91, 93)	84 (82, 85)	51 (49, 54)	28 (26, 31)
New Zealand	<40 (n=589)	99 (97, 99)	96 (94, 98)	87 (84, 90)	78 (73, 82)
	40-59 (n=1659)	97 (96, 98)	94 (92, 95)	74 (72, 77)	55 (51, 58)
	60-74 (n=1496)	96 (95, 97)	88 (87, 90)	61 (59, 64)	37 (34, 40)
	≥75 (n=342)	92 (88, 94)	83 (78, 87)	47 (41, 53)	20 (15, 25)

Figure 6.18.1**Figure 6.18.2****Table 6.9. Incident Home Dialysis Technique Survival by Age 2003-2014**

Country	Age	6 months	1 year	3 years	5 years
Australia	<40 (n=2035)	90 (88, 91)	78 (76, 80)	52 (49, 55)	39 (35, 42)
	40-59 (n=4675)	88 (87, 89)	78 (77, 80)	47 (46, 49)	29 (27, 31)
	60-74 (n=4567)	84 (83, 85)	73 (72, 75)	38 (37, 40)	19 (17, 21)
	≥75 (n=1886)	80 (78, 82)	67 (64, 69)	29 (27, 31)	10 (8, 12)
New Zealand	<40 (n=589)	89 (86, 91)	80 (76, 83)	51 (46, 56)	38 (32, 44)
	40-59 (n=1659)	89 (88, 91)	81 (79, 83)	48 (46, 51)	27 (24, 30)
	60-74 (n=1496)	88 (86, 90)	77 (74, 79)	43 (40, 45)	19 (16, 21)
	≥75 (n=342)	83 (79, 87)	71 (66, 76)	31 (26, 37)	7 (4, 11)

Figure 6.19.1




Figure 6.19.2




Table 6.10. Incident Home Dialysis Death-Censored Technique Survival by Age 2004-2015

Country	Age	6 months	1 year	3 years	5 years
Australia	<40 (n=2035)	91 (90, 92)	80 (78, 82)	56 (53, 59)	44 (41, 48)
	40-59 (n=4675)	90 (89, 91)	82 (81, 83)	57 (55, 59)	42 (40, 44)
	60-74 (n=4567)	88 (87, 89)	81 (79, 82)	57 (55, 59)	41 (39, 44)
	≥75 (n=1886)	87 (86, 89)	80 (78, 82)	59 (56, 62)	44 (39, 48)
New Zealand	<40 (n=589)	90 (88, 92)	83 (79, 86)	58 (53, 63)	48 (42, 54)
	40-59 (n=1659)	92 (90, 93)	86 (84, 88)	64 (61, 67)	47 (43, 51)
	60-74 (n=1496)	92 (90, 93)	86 (84, 88)	69 (65, 71)	52 (47, 56)
	≥75 (n=342)	91 (87, 93)	86 (81, 89)	68 (61, 74)	57 (48, 65)

Deaths on Home Dialysis

Table 6.11 shows the causes of death in patients who died while receiving home dialysis, or within 30 days of transferring from home dialysis to facility haemodialysis, during 2004-2015. Deaths from cardiovascular disease were the most common in both HHD and PD. Compared with PD patients, HHD patients were more likely to die from cardiovascular disease, but less likely to die from infection or dialysis withdrawal.

Table 6.11. Cause of Death in Home Dialysis Patients 2004-2015

Cause of death	PD	Home HD
Cardiovascular	1835 (35%)	405 (47%)
Withdrawal	1350 (26%)	141 (16%)
Cancer	233 (4%)	61 (7%)
Infection	700 (13%)	79 (9%)
Other	1079 (21%)	177 (21%)
Total	5197 (100%)	863 (100%)

Suggested Citation:

ANZDATA Registry. 39th Report, Chapter 6: Home Dialysis. Australia and New Zealand Dialysis and Transplant Registry, Adelaide, Australia. 2017. Available at: <http://www.anzdata.org.au>

ANZDATA Registry

C-1 South Australia Health and Medical Research Institute (SAHMRI)
Level 4 South
North Terrace, Adelaide
South Australia
Australia

www.anzdata.org.au
Phone +61 8 8128 4758

