

# TRANSPLANTING HOSPITAL REPORT

# 2014 - 2019

PUBLISHED December 2020 From the ANZDATA Database last surveyed on 31st December 2019



# Contents

1	Introduction	<b>2</b>
<b>2</b>	Methods	3
3	Graft Failure at 1 Year Post-transplant	<b>5</b>
	3.1 Risk-adjusted Graft Failure Ratios	5
	3.2 Funnel Plot	6
	3.3 Logistic Model Coefficients	7
4	Mortality at 1 Year Post-transplant	8
	4.1 Risk-adjusted Mortality Ratios	8
	4.2 Funnel Plot	9
	4.3 Logistic Model Coefficients	10



## 1 Introduction

This report is an abridged version of the transplant hospital report, prepared for general distribution. Individual hospital reports are also created, which contain more detailed information about the characteristics and outcomes within each hospital.

All Australian and New Zealand patients transplanted between 1 January 2014 and 31 December 2019 and who were aged  $\geq 16$  years at the time of operation were included. Both deceased and living donor transplants were included, but patients who received multi-organ grafts were excluded. Patients transplanted at a paediatric hospital were excluded. Recipients aged  $\geq 18$  years at the time of transplant were included for hospitals that provide transplantation services for all ages.

The data are based on reports to the ANZDATA Registry. Interpretation of these results must take into account both the limitations of the methodology and the context. There is considerable literature about interpretation of results from many fields, and further information can be provided for those seeking to better understand the results.

The results presented here are estimates of true values and are subject to random variation. Confidence intervals are used to present this variability. To account for the multiple comparisons made between centres, 95% false discovery rate (FDR) confidence intervals are used.

Another key limitation is the potential for factors other than those measured, which may be outside the control of treating hospitals, to affect results. This is known as residual confounding. Despite the inclusion of many factors related to patients and their care, most models predict only around 70% of the variation in transplant outcomes. ANZDATA results are consistent with international experience in this regard.

How then should results suggesting a hospital's results are inferior to expectation be interpreted? Perhaps the best approach is to consider them as signals for looking at a deeper level, bearing in mind that it may well be that the effects seen are driven by factors unrelated to the quality of care or beyond the control of individual hospitals (eg, chance, unmeasured confounders, or natural variation).

# 2 Methods

Random effects logistic regression was used to calculate the expected number of events (deaths or graft failures) at 1 year post-transplant for transplanting hospitals based on the patient and donor case-mix. Graft failure included return to dialysis or death.

Factors included in logistic regression models were recipient age, donor age, recipient gender, recipient country, presence of chronic lung disease and vascular disease (coronary artery disease, cerebrovascular disease and/or peripheral vascular disease) in the recipient, primary renal disease, number of HLA mismatches, donor type by cause of death (cerebrovascular accident (CVA), or other cause) and total ischaemic time, time since commencing RRT, peak panel reactive antibody (PRA) and transplant year. For Australian patients the peak PRA is defined as the authorised calculated PRA at match, where available, and the authorised PRA otherwise.

In addition, a random effect at the level of the transplanting hospital was included. This is a conservative assumption (i.e., this will reduce the chance of a false positive but will also reduce the chance of detecting a true positive). This term explicitly attempts to account for variation in hospital-specific performance due to chance; in doing so, it may also incorporate some elements of relevant hospital-specific outcomes. Following a similar philosophy we have included total ischaemic time in the model as a factor for adjustment; this assumption implies that this factor is not under the control of the transplanting hospital.

The logistic regression model was fitted to data from all transplant recipients to obtain the probability of an event occurring for each patient based upon their characteristics. The expected number of events was defined as the number of events expected if the patients treated at that hospital had instead been assigned at random to any hospital in Australia and New Zealand, with the random assignment weighted by hospital size. For each patient, predicted event probabilities had that patient been treated in each available hospital were calculated, then a weighted average was taken. These weighted average predicted probabilities were then summed over the patients within each hospital, resulting in the expected number of events. The standard error of the ratios of observed to expected events were estimated using 500 bootstrapped samples. Missing values for comorbidities were recoded to the comorbidity being absent. Following the comorbidities being recoded, some observations still had missing values for one or more predictor variables and these cases were excluded. We have indicated the number of missing patients for each hospital. Where this is large, then the potential exists to bias results if the pattern of missingness is not random.

For all models, the expected number of events predicted by the model within each category of covariates was consistent with the observed numbers. Additionally, the area under the receiver-operating characteristic (ROC) curves attributable to each model was also examined. For 1 year death and graft failure the area under the ROC curve were 77.0% (95% CI: 72.0-82.1) and 69.8% (95% CI: 66.3-73.2), respectively. These results indicate a model with moderate fit, and the degree of model fit is comparable with US and UK models.

This section contains the random effects logistic regression model results as odds ratios with 95% confidence intervals for the two outcomes. It also contains the number of transplants



performed by each hospital and the number of graft failure events (which includes death) and patient deaths within 1 year post-transplant. The expected number of events (graft failures or deaths) based on patient case-mix is presented as well as the ratio of observed to expected. The ratio of observed to expected is presented with a 95% false discovery rate (FDR) confidence interval, that accounts for the multiple comparisons made between centres. The expected proportion of centres identified falsely by lying outside their confidence interval is 0.05. The rate of observed events per number of transplants is also presented with a 95% confidence interval.

The ratios of observed to expected for both outcomes are also presented in funnel plots. These show the ratios on a logarithmic scale (y-axis) plotted against the effective sample size (x-axis). Hospitals with a ratio of 0 are not shown. The red line indicates a ratio of 1, and the contours indicate 95% FDR confidence intervals. If a hospital lies within the confidence intervals then that hospital has an observed to expected ratio that is statistically consistent (at a 5% FDR level) with 1 (i.e. there is no statistical difference in the number of observed and expected events). If a hospital lies above the upper control lines, this indicates that the number of observed deaths is statistically greater than the number expected under the model. Conversely, if a hospital lies below the lines, this indicates statistically fewer observed deaths than expected under the model. The SMR is presented on a logarithmic scale as confidence intervals for the logarithm of the SMR (log-SMR) have better coverage properties. The effective sample size measures the variability of each log-SMR relative to the overall variability of all log-SMRs.

In interpreting the SMR and funnel plots it should be borne in mind that the precision of these estimates is strongly influenced by the number of patients in a hospital. As such, smaller hospitals will have less precise estimates and greater uncertainty about where the true effect lies. This is shown in wider confidence intervals for the SMR estimates and likely greater change in these estimates as they are updated over time.

### Graft Failure at 1 Year Post-transplant 3

#### **Risk-adjusted Graft Failure Ratios** 3.1

Code	Hospital	$\operatorname{Grafts}^*$	Observed	Expected	Observed Rate (95% CI)	Ratio (95% FDR CI)
ALFD	Alfred Hospital	141(9)	7	6.2	0.050 (0.020-0.100)	1.13 (0.27-4.67)
AUCK	Auckland Hospital	469(0)	13	19.3	$0.028\ (0.015 - 0.047)$	0.67(0.30-1.49)
AUST	Austin Hospital	284(25)	11	10.8	$0.039\ (0.019 - 0.068)$	1.02(0.39-2.66)
CHCH	Christchurch Hospital	133(2)	5	4.3	$0.038\ (0.012 - 0.086)$	1.16 (0.27-4.90)
CNAR	Central Northern Adelaide Renal Service	393(13)	18	15.6	$0.053 \ (0.033 - 0.081)$	1.15(0.59-2.25)
FSTH	Fiona Stanley Hospital	162(18)	4	4.7	$0.025\ (0.007-0.062)$	$0.86\ (0.16-4.52)$
HUNT	John Hunter Hospital	186(15)	9	7.4	0.070(0.038 - 0.117)	1.22(0.45-3.28)
MMCA	Monash Medical Centre (Adults)	377 (17)	21	17.0	$0.066\ (0.043-0.096)$	1.24(0.66-2.31)
POWH	Prince Of Wales Hospital	206(1)	7	9.4	$0.034 \ (0.014 - 0.069)$	0.75(0.23-2.43)
QRTS	Queensland Kidney Transplant Service	799 (14)	24	29.2	$0.030\ (0.019-0.044)$	0.82(0.46-1.46)
RLPT	Royal Perth Hospital	30(12)	0	0.8	$0.000 \ (0.000-0.116)$	0.00 ()
RMBH	Royal Melbourne Hospital	646(50)	22	24.6	$0.037 \ (0.024 - 0.055)$	0.89(0.48-1.65)
RNSH	Royal North Shore Hospital	174(0)	7	7.3	$0.040\ (0.016-0.081)$	0.95(0.32-2.86)
SCGH	Sir Charles Gairdner Hospital	256(39)	13	8.2	$0.059\ (0.033-0.095)$	1.58(0.69-3.62)
STVI	St Vincent's Hospital (NSW)	18(4)	2	0.5	0.167(0.036-0.414)	4.05 (0.75-21.98)
STVP	St Vincent's Private Hospital (NSW)	18(0)	0	0.4	0.000(0.000-0.185)	0.00 ()
SVIN	St Vincent's Hospital (VIC)	125(25)	2	5.0	0.040(0.013-0.091)	$0.40\ (0.05-3.13)$
SWRS	Statewide Renal Services	543 (66)	27	20.1	$0.053 \ (0.036 - 0.076)$	1.34 (0.80-2.25)
WELN	Wellington Hospital	167 (11)	10	5.0	0.060 (0.029-0.107)	2.00 (0.95-4.20)
WEST	Western Renal Service	362 (33)	10	15.2	0.033 (0.017 - 0.057)	0.66 (0.25 - 1.77)

Table 1: Observed vs expected number of graft failures at 1 year post transplant

СЛ

Transplant procedures performed 2014-2018 \* The number in brackets is the number of grafts excluded from logistic regression due to missing data

### 3.2 Funnel Plot



Missing comorbidities are recoded to being absent Observations with other missing values are dropped from the model

6

## 3.3 Logistic Model Coefficients

Table 2: Logistic regression model odds ratios for graft failure at 1 year post transplant

	Odds Ratio	95% CI
Subsequent Graft	0.96	[0.57, 1.60]
Age (years)	1.01	[1.00, 1.02]
Male	1.66	[1.19, 2.30]
Duration of ESKD		
Preemptive	0.65	[0.30, 1.40]
< 6  months	0.42	[0.17, 1.06]
6-12 months	0.68	[0.37, 1.24]
12-23 months	0.68	[0.44, 1.05]
24 + months	ref.	
Peak PRA	1.01	[1.00, 1.01]
Donor Age (years)		
<15 years	1.91	[0.84, 4.33]
15-49 years	ref.	
50-59 years	1.95	[1.36, 2.79]
$\geq 60$ years	1.73	[1.18, 2.54]
New Zealand	1.10	[0.67, 1.78]
Lung Disease	1.05	[0.67, 1.64]
Cardiac Disease	1.47	[1.07, 2.01]
PRD		
Glomerulonephritis	ref.	
Polycystic Disease	0.91	[0.54, 1.53]
Diabetic Nephropathy	1.47	[0.97, 2.21]
Miscellaneous	1.37	[0.97, 1.92]
Number of Mismatches	1.01	[0.93, 1.11]
Year of Transplant		
2014	ref.	
2015	1.13	[0.73, 1.75]
2016	0.86	[0.55, 1.36]
2017	0.68	[0.42, 1.08]
2018	0.76	[0.48, 1.19]
Donor Type / Cause of Death / Ischaemic Time		L , J
LD	ref.	
DD CVA 0-11 hours	2.01	[1.17, 3.43]
DD non-CVA 0-11 hours	1.61	[0.93, 2.79]
DD CVA $>12$ hours	1.76	[1.01, 3.08]
DD non- $\overline{CVA} > 12$ hours	1.68	[0.96.2.95]

### Mortality at 1 Year Post-transplant 4

#### **Risk-adjusted Mortality Ratios** 4.1

 $\infty$ 

Code	Hospital	Patients <sup>*</sup>	Observed	Expected	Observed Rate (95% CI)	Ratio (95% FDR CI)
ALFD	Alfred Hospital	122 (7)	1	2.0	$0.008 \ (0.000 - 0.045)$	0.51 (0.11-2.39)
AUCK	Auckland Hospital	427 (0)	9	8.9	0.021 (0.010-0.040)	1.01 (0.47-2.16)
AUST	Austin Hospital	259 (22)	4	4.8	$0.015\ (0.004-0.039)$	0.84 (0.21-3.33)
CHCH	Christchurch Hospital	122(2)	2	1.7	$0.016\ (0.002-0.058)$	1.15 (0.25-5.22)
CNAR	Central Northern Adelaide Renal Service	341 (11)	8	5.9	$0.026\ (0.012 - 0.050)$	1.35(0.50-3.62)
FSTH	Fiona Stanley Hospital	136(16)	0	1.4	0.000 (0.000-0.027)	0.00 ()
HUNT	John Hunter Hospital	162(14)	4	2.8	$0.037 \ (0.014 - 0.079)$	1.42(0.33-6.19)
MMCA	Monash Medical Centre (Adults)	325(13)	9	6.4	$0.028 \ (0.013 - 0.052)$	1.42(0.59-3.37)
POWH	Prince Of Wales Hospital	190(1)	4	4.1	$0.021 \ (0.006 - 0.053)$	0.98 (0.24-4.03)
QRTS	Queensland Kidney Transplant Service	693(14)	4	11.1	$0.006 \ (0.002 - 0.015)$	$0.36\ (0.09-1.45)$
RLPT	Royal Perth Hospital	24(11)	0	0.1	$0.000 \ (0.000 - 0.142)$	0.00 ()
RMBH	Royal Melbourne Hospital	520 (37)	8	8.7	$0.017 \ (0.008 - 0.033)$	0.92(0.36-2.31)
RNSH	Royal North Shore Hospital	152(0)	5	3.2	$0.033\ (0.011 - 0.075)$	1.57 (0.48-5.11)
SCGH	Sir Charles Gairdner Hospital	217(35)	5	3.0	$0.023 \ (0.008 - 0.053)$	1.68 (0.42-6.70)
STVI	St Vincent's Hospital (NSW)	15(3)	0	0.1	$0.000 \ (0.000-0.218)$	0.00 ()
STVP	St Vincent's Private Hospital (NSW)	18(0)	0	0.1	$0.000 \ (0.000 - 0.185)$	0.00 ()
SVIN	St Vincent's Hospital (VIC)	111(22)	0	2.0	$0.009\ (0.000-0.049)$	0.00 ()
SWRS	Statewide Renal Services	490(59)	11	8.0	$0.027 \ (0.014 - 0.045)$	1.38 (0.61-3.08)
WELN	Wellington Hospital	155(10)	2	2.2	$0.013 \ (0.002 - 0.046)$	0.91 (0.20-4.15)
WEST	Western Renal Service	316(29)	7	5.8	0.022 (0.009 - 0.045)	1.21 (0.43-3.38)

Table 3: Observed vs expected number of deaths at 1 year post transplant

Patients transplanted 2014-2018 (primary grafts only)
<sup>\*</sup> The number in brackets is the number of patients excluded from logistic regression due to missing data

### 4.2 Funnel Plot



Missing comorbidities are recoded to being absent Observations with other missing values are dropped from the model

## 4.3 Logistic Model Coefficients

Table 4: Logistic regression model odds ratios for mortality at 1 year post transplant

	Odds Ratio	95% CI
Age (years)	1.04	[1.02, 1.07]
Male	1.40	[0.82, 2.38]
Duration of ESKD		
Preemptive	0.77	[0.22, 2.65]
< 6  months	0.46	[0.11, 1.98]
6-12 months	0.91	[0.40, 2.10]
12-23 months	0.52	[0.25, 1.07]
24+ months	ref.	
Peak PRA	1.01	[1.00, 1.02]
Donor Age (years)		
<15 years	1.85	[0.53, 6.41]
15-49 years	ref.	
50-59 years	1.26	[0.70, 2.28]
$\geq 60$ years	1.53	[0.86, 2.71]
New Zealand	1.44	[0.75, 2.80]
Lung Disease	1.58	[0.86, 2.91]
Cardiac Disease	1.88	[1.16, 3.05]
PRD		
Glomerulonephritis	ref.	
Polycystic Disease	1.35	[0.63, 2.88]
Diabetic Nephropathy	1.53	[0.82, 2.87]
Miscellaneous	1.46	[0.81, 2.64]
Number of Mismatches	1.03	[0.89, 1.18]
Year of Transplant		
2014	ref.	
2015	1.63	[0.78, 3.39]
2016	1.16	[0.54, 2.50]
2017	1.13	[0.53, 2.40]
2018	0.90	[0.42, 1.95]
Donor Type / Cause of Death / Ischaemic Time		
LD	ref.	
DD CVA 0-11 hours	2.57	[1.01, 6.51]
DD non-CVA 0-11 hours	2.04	[0.79, 5.28]
DD CVA $\geq 12$ hours	1.06	[0.36, 3.14]
DD non-CVA $\geq 12$ hours	3.06	[1.21, 7.76]